



Contribution ID: 40

Type: **Oral**

## On the direct correlation between gamma-rays and PeV neutrinos from blazars

*Friday, December 9, 2016 2:40 PM (20 minutes)*

We study the frequently used assumption in multi-messenger astrophysics that the gamma-ray and neutrino fluxes are directly connected because they are assumed to be produced by the same photohadronic production chain. An interesting candidate source for this test is the flat-spectrum radio quasar PKS B1424-418, which recently called attention of a potential correlation between an IceCube PeV-neutrino event and its burst phase. We simulate both the multi-waveband photon and the neutrino emission from this source using a self-consistent radiation model. We demonstrate that a simple hadronic model cannot adequately describe the spectral energy distribution for this source, but a lepto-hadronic model with sub-dominant hadronic component can reproduce the multi-waveband photon spectrum observed during various activity phases of the blazar. As a conclusion, up to about 0.3 neutrino events may coincide with the burst, which implies that the leptonic contribution dominates in the relevant energy band. We also demonstrate that the time-wise correlation between the neutrino event and burst phase is weak.

**Author:** Dr GAO, Shan (DESY)

**Co-authors:** Prof. POHL, Martin (DESY and U Potsdam); Prof. WINTER, Walter (DESY)

**Presenter:** Dr GAO, Shan (DESY)

**Session Classification:** MWL and Multi-Messenger

**Track Classification:** HAP Workshop