

Flavorful Ways to New Physics


Waldhotel Zollernblick, Oct 28-31, 2014



Flavor Physics at Belle and Belle II


Peter Križan

University of Ljubljana and J. Stefan Institute



University
of Ljubljana

"Jožef Stefan"
Institute






Contents

- Introduction with a little bit of B factory primer
- B factories: recent results
- Super B factory: status and outlook
- Summary



Flavour physics at the luminosity frontier with asymmetric B factories




<p>BaBar</p> <p>Belle</p>	<p>$p(e^-) = 9 \text{ GeV}$</p> <p>$p(e^-) = 8 \text{ GeV}$</p>	<p>$p(e^+) = 3.1 \text{ GeV}$</p> <p>$p(e^+) = 3.5 \text{ GeV}$</p>
---	---	---

$\beta\gamma = 0.56$
$\beta\gamma = 0.42$


To a large degree shaped flavour physics in the previous decade

Advantages of B factories in the LHC era




Unique capabilities of B factories:

- Exactly two B mesons produced (at $\Upsilon(4S)$)
- High flavour tagging efficiency
- Detection of gammas, π^0 s, K_L s
- Very clean detector environment (can observe decays with several neutrinos in the final state!)
- Well understood apparatus, with known systematics, checked on control channels



Integrated luminosity at B factories



> 1 ab⁻¹

On resonance:

$\Upsilon(5S)$: 121 fb⁻¹

$\Upsilon(4S)$: 711 fb⁻¹

$\Upsilon(3S)$: 3 fb⁻¹

$\Upsilon(2S)$: 25 fb⁻¹

$\Upsilon(1S)$: 6 fb⁻¹

Off reson./scan:

~ 100 fb⁻¹

~ 550 fb⁻¹

On resonance:

$\Upsilon(4S)$: 433 fb⁻¹

$\Upsilon(3S)$: 30 fb⁻¹


$\Upsilon(2S)$: 14 fb⁻¹

Off resonance:

~ 54 fb⁻¹




CP violation in the B system and unitarity triangle



B factories: CP violation in the B system

CP violation in the B system: from the **discovery** (2001) to a **precision measurement** (2011).



Comparison of **energy** / **intensity** frontiers

To observe a large ship far away one can either use **strong binoculars** or observe **carefully the direction and the speed of waves** produced by the vessel.

Energy frontier (LHC)




**Luminosity frontier -
(super) B factories**

The unitarity triangle – new/final measurements

Constraints from measurements of angles and sides of the unitarity triangle → Remarkable agreement, but still 10-20% NP allowed

Selected results:


- $\sin 2\phi_1 (= \sin 2\beta)$: final measurements
- $\phi_2 (= \alpha)$: final measurements
- $\phi_3 (= \gamma)$: new model-independ. method
- Rare decays



CP violation measurement

Want to measure the asymmetry between B and anti-B mesons,

$$P(B^0(\bar{B}^0) \rightarrow f_{CP}, t) = e^{-\Gamma t} (1 \mp \sin(2\phi_1) \sin(\Delta m t))$$




→ Want to distinguish the decay rate of B (dotted) from the decay rate of anti-B (full).

Integrals are equal, time information mandatory! (true at $Y(4s)$, but not for incoherent production)

Resolution \sim B lifetime

B meson production at $\Upsilon(4S)$




Cross Sections at $\Upsilon(4S)$:

$b\bar{b} \sim 1.1 \text{ nb}$

$c\bar{c} \sim 1.3 \text{ nb}$

$d\bar{d}, s\bar{s} \sim 0.3 \text{ nb}$


$u\bar{u} \sim 1.4 \text{ nb}$



$e^+e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B}$
 $L = 1$ state

CP violation measurement


Measure the difference in time evolution in B^0 and anti- B^0 decays to a CP eigenstate




CMS should be boosted!

Experimental considerations


Detector form: symmetric for symmetric energy beams; **slightly extended in the boost direction** for an asymmetric collider.



Exaggerated plot: in reality $\beta\gamma=0.5$




CLEO





BELLE


Belle spectrometer at KEK-B



Reconstruction of rare B meson decays




Reconstructing rare B meson decays at Y(4s): use two variables,
beam constrained mass M_{bc}
 and
energy difference ΔE



$$\Delta E \equiv \sum E_i - E_{CM}/2$$

$$M_{bc} = \sqrt{(E_{CM}/2)^2 - (\sum \vec{p}_i)^2}$$


Continuum suppression



$e^+e^- \rightarrow qq$ "continuum" ($\sim 3x$ BB)


To suppress: use event shape variables

Continuum
Jet-like



BB


spherical



CP violation measurement

Want to measure the asymmetry between B and anti-B mesons,

$$P(B^0(\bar{B}^0) \rightarrow f_{CP}, t) = e^{-\Gamma t} (1 \mp \sin(2\phi_1) \sin(\Delta m t))$$



→ Want to distinguish the decay rate of B (dotted) from the decay rate of anti-B (full).


Integrals are equal, time information mandatory! (true at Y(4s), but not for incoherent production)

Resolution \sim B lifetime





Final measurement of $\sin 2\phi_1 (= \sin 2\beta)$

ϕ_1 from CP violation measurements in $B^0 \rightarrow c\bar{c} K^0$



Final measurement: with improved tracking, more data, improved systematics (50% more statistics than last result with 492 fb^{-1});
 $cc = J/\psi, \psi(2S), \chi_{c1} \rightarrow 25\text{k events}$

Detector effects: wrong tagging, finite Δt resolution \rightarrow determined using control data samples



Belle, final, 710 fb^{-1} , PRL 108, 171802 (2012)

K_L detection


Important cross check:

Measure CP violation for $B \rightarrow$
CP=+1 eigenstate

$\rightarrow B \rightarrow J/\psi K_L$


Need a detector for K_L s – muon
detections system acts as a
hadron calorimeter

Measure only the K_L interaction
point coordinate, not the K_L
energy.





Final measurements of $\sin 2\phi_1 (= \sin 2\beta)$



ϕ_1 from $B^0 \rightarrow c\bar{c} K^0$

Final results for $\sin 2\phi_1$

Belle: $0.668 \pm 0.023 \pm 0.012$
 BaBar: $0.687 \pm 0.028 \pm 0.012$

Belle, PRL 108, 171802 (2012)

BaBar, PRD 79, 072009 (2009)




with a single experiment precision of $\sim 4\%$!

Comparison with LHCb:

- The power of tagging at B factories: **33%** vs $\sim 2\text{-}3\%$ at LHCb
- LHCb: with 8k tagged $B_d \rightarrow J/\psi K_S$ events from 1/fb measured $\sin 2\beta = 0.73 \pm 0.07(\text{stat.}) \pm 0.04(\text{syst.})$
- Uncertainties at B factories - e.g., Belle final result $\sin 2\beta = 0.668 \pm 0.023(\text{stat.}) \pm 0.012(\text{syst.})$ - are **3x smaller** than at LHCb

Final measurement of $\phi_2 (\alpha)$ in $B \rightarrow \pi^+\pi^-$ decays

ϕ_2 from CP violation measurements in $B^0 \rightarrow \pi^+\pi^-$



Belle, 710 fb⁻¹
PRD **88**, 092003 (2013)

$$a_{f_{CP}} = C \cos(\Delta mt) + S \sin(\Delta mt)$$



Belle, this measurement:

$$S = -0.64 \pm 0.08 \pm 0.03$$

$$C = -0.33 \pm 0.06 \pm 0.03$$

BaBar:

$$S = -0.68 \pm 0.10 \pm 0.03$$

$$C = -0.25 \pm 0.08 \pm 0.02$$



Measurement of $B \rightarrow \pi^0\pi^0$ decays


ϕ_2 from CP violation measurements in $B^0 \rightarrow \pi^+\pi^-$
Extraction not easy because of the penguin contribution

BR for the $B \rightarrow \pi^0\pi^0$ decay important to resolve this issue.

Hard channel to measure: four gammas, continuum (ee \rightarrow qq) background

- Theory: BR < 1x10⁻⁶ (Phys.Rev.D83:034023,2011)
- Belle, **1/3 of data** PRL 94, 181803(2005) = (2.32 +0.4-0.5 +0.2-0.3) 10⁻⁶
- BaBar PR D87 052009 (1.83 \pm 0.21 \pm 0.13) 10⁻⁶





Belle new result with full data set: Improved rejection of out-of-time electromagnetic calorimeter hits (some of which contribute to a peaking background).



How the penguin distorts the tree level measurement

Pit Vanhoefer, CKM2014

Measurement of $B \rightarrow \pi^0\pi^0$ decays



Preliminary

$$Br(B \rightarrow \pi^0\pi^0) = (0.90 \pm 0.20 \text{ (stat)} \pm 0.15 \text{ (syst)}) \cdot 10^{-6}$$


(6.7 σ significance)

A_{CP} under preparation \rightarrow stay tuned



Improved measurement of $\phi_2 (\alpha)$ in $B \rightarrow \pi\pi, \rho\rho, \rho\pi$ decays


$\phi_2 (\alpha)$ from CP violation and branching fraction measurements in $B \rightarrow \pi\pi, \rho\rho, \rho\pi$



$\phi_2 = \alpha = (85.4^{+4.0}_{-3.8})$ degrees

http://ckmfitter.in2p3.fr/www/results/plots_fpcp13/ckm_res_fpcp13.html


Still to be updated for the final version: new results expected from Belle on $\rho^+\rho^-, \rho\pi$; a new $\rho\pi$, analysis published by BaBar \rightarrow PRD88, 012003 (2013).




$\phi_3 (= \gamma)$ with Dalitz analysis

GGSZ method:

The best way to measure ϕ_3



A. Giri et al., PRD68, 054018 (2003)
 A. Bondar et al (Belle), Proc. BINP Meeting on Dalitz Analyses, 2002



$$\bar{D}^0 \rightarrow K_S \pi^+ \pi^-$$

3-body $D^0 \rightarrow K_S \pi^+ \pi^-$ Dalitz amplitude

$$|M_{\pm}(m_+^2, m_-^2)|^2 = |f_D(m_+^2, m_-^2) + re^{i\delta_B \pm i\phi_3} f_D(m_-^2, m_+^2)|^2$$

Model dependent description of f_D using continuum D^* data \Rightarrow systematic uncertainty

$$= \left| \left[\text{Dalitz Plot 1} \right] + re^{i\delta_B \pm i\phi_3} \left[\text{Dalitz Plot 2} \right] \right|^2$$

$$\phi_3 = (78 \pm 12 \pm 4 \pm 9)^\circ$$

$$\phi_3 = (68 \pm 14 \pm 4 \pm 3)^\circ$$

Belle, PRD81, 112002, (2010), 605 fb⁻¹

BaBar, PRL 105, 121801, (2010)



$\phi_3 (= \gamma)$ from model-independent/binning Dalitz method

GGSZ method: How to avoid the model dependence?

→ Suitably subdivide the Dalitz space into bins

$$M_i^\pm = h \{ K_i + r_B^2 K_{-i} + 2\sqrt{K_i K_{-i}} (x_\pm c_i + y_\pm s_i) \}$$

$$x_\pm = r_B \cos(\delta_B \pm \phi_3) \quad y_\pm = r_B \sin(\delta_B \pm \phi_3)$$



M_i : # B decays in bins of D Dalitz plane, K_i : # D^0 (\bar{D}^0) decays in bins of D Dalitz plane ($D^* \rightarrow D\pi$), c_i, s_i : strong ph. difference between symm. Dalitz points ← Cleo, PRD82, 112006 (2010)



Use only DK
 $N_{sig} = 1176 \pm 43$


Belle, 710 fb⁻¹, Phys. Rev. D85 (2012) 112014

4-dim fit for signal yield
($\Delta E, M_{bc}, \cos\theta_{thrust}, \mathcal{F}$);

$$\phi_3 = (77.3 \pm 15 \pm 4.1 \pm 4.3)^\circ$$

from c_i, s_i (statist.!) →

to be reduced with BESIII data →



Peter Križan, Ljubljana


New method pioneered by Belle, very important for large event samples at LHCb and super B factory

ϕ_3 measurement

Combined ϕ_3 value:





$$\phi_3 = (67 \pm 11) \text{ degrees}$$

Note that at B factories the measurement of ϕ_3 finally turned out to be much better than expected!



This is not the last word from B factories, analyses still to be finalized...





Rare B decays



CP violation in penguin dominated $b \rightarrow$ qqs transitions

CP violation given by the same parameter $\sin 2\phi_1$ as in $J/\psi K$ decays


\rightarrow to be published in JHEP



Decay mode	$-\xi_f \mathcal{S}_f$	\mathcal{A}_f
$\eta' K_S^0$	$+0.71 \pm 0.07$	$+0.02 \pm 0.05$
$\eta' K_L^0$	$+0.46 \pm 0.21$	$+0.09 \pm 0.14$
$\eta' K^0$	$+0.68 \pm 0.07 \pm 0.03$	$+0.03 \pm 0.05 \pm 0.04$

Advantage of B factories!


$B \rightarrow X_s \gamma$ inclusive



Radiative decay sensitive to charged Higgs

- continuum
- $\pi^0 \rightarrow \gamma\gamma$
- $\eta \rightarrow \gamma\gamma$
- $b \rightarrow s\gamma$


Experimentally difficult \dashrightarrow



Experiment: measure low E_γ
 \Rightarrow huge bkg. $\Rightarrow E_\gamma > E_{cut}$

Theory:
 parameter extraction from
 partial $\text{Br}(E_\gamma > E_{cut}) \Rightarrow$
 extrapolation needed;

Only γ on signal side reconstructed
 Improve S/B by tagging the other B



$$\mathcal{B}(B \rightarrow X_s \gamma; 1.7 \text{ GeV} < E_\gamma < 2.8 \text{ GeV}) = (3.47 \pm 0.15 \pm 0.40) \cdot 10^{-4}$$

B \rightarrow X_sγ inclusive


Branching fraction,
world average

$$\mathcal{B}(B \rightarrow X_s \gamma; E_\gamma > 1.6 \text{ GeV}) = (3.55 \pm 0.24(\text{stat.} + \text{syst.}) \pm 0.09(\text{shape } f.) \cdot 10^{-4}$$

HFAG, ICHEP'10

Decay rate sensitive to
charged Higgs

→ tight constraints on models of new physics, two-Higgs-doublet model II mass limit at $\sim 300 \text{ GeV}/c^2$



Measurements systematics dominated

Systematics can be reduced by stronger tagging (e.g. full reconstruction of the other B) on the account of stat. uncertainty


⇒ need a larger sample → Super B factory

B \rightarrow $X_s \gamma$, semi-inclusive



Sum of 38 exclusive channels

Mode ID	Final State	Mode ID	Final State
1	$K^+ \pi^-$	20	$K_S^0 \pi^+ \pi^0 \pi^0$
2	$K_S^0 \pi^+$	21	$K^+ \pi^+ \pi^- \pi^0 \pi^0$
3	$K^+ \pi^0$	22	$K_S^0 \pi^+ \pi^- \pi^0 \pi^0$
4	$K_S^0 \pi^0$	23	$K^+ \eta$
5	$K^+ \pi^+ \pi^-$	24	$K_S^0 \eta$
6	$K_S^0 \pi^+ \pi^-$	25	$K^+ \eta \pi^-$
7	$K^+ \pi^+ \pi^0$	26	$K_S^0 \eta \pi^+$
8	$K_S^0 \pi^+ \pi^0$	27	$K^+ \eta \pi^0$
9	$K^+ \pi^+ \pi^- \pi^-$	28	$K_S^0 \eta \pi^0$
10	$K_S^0 \pi^+ \pi^+ \pi^-$	29	$K^+ \eta \pi^+ \pi^-$
11	$K^+ \pi^+ \pi^- \pi^0$	30	$K_S^0 \eta \pi^+ \pi^-$
12	$K_S^0 \pi^+ \pi^- \pi^0$	31	$K^+ \eta \pi^- \pi^0$
13	$K^+ \pi^+ \pi^+ \pi^- \pi^-$	32	$K_S^0 \eta \pi^+ \pi^0$
14	$K_S^0 \pi^+ \pi^+ \pi^- \pi^-$	33	$K^+ K^+ K^-$
15	$K^+ \pi^+ \pi^- \pi^- \pi^0$	34	$K^+ K^- K_S^0$
16	$K_S^0 \pi^+ \pi^+ \pi^- \pi^0$	35	$K^+ K^+ K^- \pi^-$
17	$K^+ \pi^0 \pi^0$	36	$K^+ K^- K_S^0 \pi^+$
18	$K_S^0 \pi^0 \pi^0$	37	$K^+ K^+ K^- \pi^0$
19	$K^+ \pi^- \pi^0 \pi^0$	38	$K^+ K^+ K_S^0 \pi^0$




Branching fraction,
(corresponding to a minimum photon
energy of 1.9 GeV)



$$\mathcal{B}(B \rightarrow X_s \gamma; M_{X_s} < 2.8 \text{ GeV} / c^2) =$$

$$= (3.51 \pm 0.17(\text{stat.}) \pm 0.33(\text{syst})) \cdot 10^{-4}$$

To be submitted to PRD




Example of a missing energy decay




Full reconstruction tagging

Idea: **fully reconstruct** one of the B's to tag B flavor/charge, determine its momentum, and exclude decay products of this B from further analysis
(exactly two B's produced in $\Upsilon(4S)$ decays)



Powerful tool for B decays with neutrinos, used in several analyses in this talk

→ unique feature at B factories



Method: tag one B with full reconstruction, look for the $B^- \rightarrow \tau^- \nu_\tau$ in the rest of the event.

Main discriminating variable on the signal side:
remaining energy in the calorimeter, not associated with any charged track or photon
 → Signal at $E_{ECL} = 0$



Belle $Br(B \rightarrow \tau \nu) = [0.72^{+0.27}_{-0.25} \pm 0.11] \times 10^{-4}$
 PRL 110, 131801 (2013)

BaBar $Br(B \rightarrow \tau \nu) = [1.83^{+0.53}_{-0.49} \pm 0.24] \times 10^{-4}$
 Phys. Rev. D 88, 031102(R) (2013)


All measurements combined

$$BF(B \rightarrow \tau \nu) = (1.15 \pm 0.23) \cdot 10^{-4}$$

$$r_H = \frac{BF(B \rightarrow \tau \nu)_{meas}}{BF(B \rightarrow \tau \nu)_{SM}} = 1.14 \pm 0.40$$



Charged Higgs limits from $B \rightarrow \tau^- \nu_\tau$




$$r_H = \frac{BF(B \rightarrow \tau \nu)}{BF(B \rightarrow \tau \nu)_{SM}} = \left(1 - \frac{m_B^2}{m_H^2} \tan^2 \beta \right)^2$$


Measured value

→ limit on charged Higgs mass vs. $\tan\beta$
(for type II 2HDM)

B factories: Exclusion plot




Super B factory: Discovery plot: very much competitive with LHC!



Belle update $B^- \rightarrow \tau^- \nu_\tau$

Method: tag with a semileptonic B decay, look for the $B^- \rightarrow \tau^- \nu_\tau$ in the rest of the event.

Again: Main discriminating variable on the signal side: **remaining energy in the calorimeter**, not associated with any charged track or photon
→ Signal at $E_{ECL} = 0$




Belle update $B^- \rightarrow \tau^- \nu_{\tau}$ tag with a semileptonic B

Belle-CONF-1401

- $\mathcal{B}(B \rightarrow \tau \nu) = [1.25 \pm 0.28 \text{ (stat)} \pm 0.27 \text{ (syst)}] \times 10^{-4}$
- Signal significance of 3.4σ including systematics

Decay Mode	N_{sig}	$\mathcal{B}(10^{-4})$
$\tau^- \rightarrow \mu^- \nu_{\tau} \bar{\nu}_{\mu}$	13 ± 21	0.34 ± 0.55
$\tau^- \rightarrow e^- \nu_{\tau} \bar{\nu}_e$	47 ± 25	0.90 ± 0.47
$\tau^- \rightarrow \pi^- \nu_{\tau}$	57 ± 21	1.82 ± 0.68
$\tau^- \rightarrow \rho^- \nu_{\tau}$	119 ± 33	2.16 ± 0.60
Combined	222 ± 50	1.25 ± 0.28

statistical errors only




- Consistent results among tau channels

➔ Central value shifted towards SM

- Combination with Belle hadronic tag result in progress

B → D^(*)τν decays

Semileptonic decay sensitive to charged Higgs



Ratio of τ to μ,e could be reduced/enhanced significantly
Kamenik, Mescia arXiv:0802.3790

$$R(D) \equiv \frac{\mathcal{B}(B \rightarrow D\tau\nu)}{\mathcal{B}(B \rightarrow D\ell\nu)}$$

Complementary and competitive with B → τν


1. Smaller theoretical uncertainty of R(D)

(For B → τν,
There is O(10%) f_B uncertainty from lattice QCD)

2. Large Brs (~1%) in SM (Ulrich Nierste arXiv:0801.4938.)

3. Differential distributions can be used to discriminate W⁺ and H⁺

4. Sensitive to different vertex B → τ ν: H-b-u, B → Dτν: H-b-c
(LHC experiments sensitive to H-b-t)



First observation of B → D^{*-}τν by Belle (2007)

→ PRL 99, 191807 (2007)

$B \rightarrow D^{(*)} \tau \nu$ decays

Exclusive hadron tag data



$$\mathcal{R}(D)_{\text{exp}} = 0.440 \pm 0.072 \quad \mathcal{R}(D^*)_{\text{exp}} = 0.332 \pm 0.030$$


$$\updownarrow 2.0\sigma$$

$$\updownarrow 2.7\sigma$$


$$\mathcal{R}(D)_{\text{SM}} = 0.297 \pm 0.017 \quad \mathcal{R}(D^*)_{\text{SM}} = 0.252 \pm 0.003$$

SM expectations in S. Fajfer, J. Kamenik, I. Nisandzic, PRD 85, 094025 (2012).

→ Combined result: 3σ away from SM.



Blue: this result, red: Type-II 2HDM.





→ Combined result: Type II 2HDM excluded at 99.8% C.L. for any values of $\tan\beta$ and charged Higgs mass

B \rightarrow $h\nu\bar{\nu}$ decays

Method: again tag one B with full reconstruction, search for signal in the remaining energy in the calorimeter, at $E_{ECL} = 0$

Present status: recent update from Belle



$$N_{Sig} = 13.3_{-6.6}^{+7.4} (stat) \pm 2.3 (syst)$$

$$S_{stat+syst} = 2.0\sigma$$


Belle, Phys. Rev. D 87, 111103(R) (2013)

$B \rightarrow K^{(*)} \nu \bar{\nu}$

arXiv:1002.5012

adopted from W. Altmannshofer et al.,
JHEP 0904, 022 (2009)




SM: penguin + box diagrams



$$B \rightarrow K \nu \bar{\nu}, \mathcal{B} \sim 4 \cdot 10^{-6}$$

$$B \rightarrow K^* \nu \bar{\nu}, \mathcal{B} \sim 6.8 \cdot 10^{-6}$$

Look for deviations from the expected values \rightarrow information on anomalous couplings C_{R}^{ν} and C_{L}^{ν} compared to $(C_{L}^{\nu})^{\text{SM}}$



Charm and τ physics

B factories = charm and τ factories

Charm and τ can be found in any "Y(nS) samples"

→ the integrated luminosity of the samples used for charm and τ studies is larger than for the B physics studies (Belle $\sim 1 \text{ ab}^{-1}$, BaBar $\sim 0.550 \text{ ab}^{-1}$)

→ This will of course remain true for the super B factory

A few examples of the strengths of B factories:

- CP violation in charm at B factories (and super B factories) → can measure CPV **separately** in individual decay channels, $\pi^+\pi^-$, K^+K^- , $K_S \pi$, ...
- $D\bar{D}$ pairs produced with **very few** light hadrons
- **Full reconstruction** of events



Rare charm decays: tag with the other D

Again make use of the **hermeticity of the apparatus!**

Example: leptonic decays of D_s

$$e^+ e^- \rightarrow c\bar{c} \rightarrow \bar{D}_{\text{tag}} K X_{\text{frag}} D_s^{*+}$$

Recoil method in charm events:

- Reconstruct D_{tag} to tag charm, kaon to tag strangeness
- Additional light mesons (X_{frag}) can be produced in the fragmentation process ($\pi, \pi\pi, \dots$)

2 step reconstruction:

- Inclusive reconstruction of D_s mesons for normalization (without any requirements upon D_s decay products)
- Within the inclusive D_s sample search for D_s decays
 - $D_s \rightarrow \mu\nu$: peak at $m_\nu^2 = 0$ in $M_{\text{miss}}^2(D_{\text{tag}} K X_{\text{frag}} \gamma \mu)$
 - $D_s \rightarrow \tau\nu$: peak towards 0 in extra energy in calorimeter

$$D_s^+ \rightarrow \mu^+ \nu_\mu$$




Fit to the missing mass squared – $M_{\text{miss}}^2(D_{\text{tag}} K X_{\text{frag}} \gamma \mu^\pm)$

$$D_s^+ \rightarrow \mu^+ \nu_\mu$$

Belle Preliminary (913 fb⁻¹)

Selection:

- $M_{\text{miss}}(D_{\text{tag}} K X_{\text{frag}} \gamma)$ signal region
- 1 charged track pointing to the IP
- passing muon PID requirements



$$N_{D_s \rightarrow \mu \nu}^{\text{excl}} = 489 \pm 26$$

Belle, arxiv:1301.7218

Belle preliminary @ 913 fb⁻¹

$$\mathcal{B}(D_s^+ \rightarrow \mu^+ \nu_\mu) = (0.528 \pm 0.028(\text{stat.}) \pm 0.019(\text{syst.}))\%$$

Most precise measurement up to date.




Extract f_{D_s} :

$$f_{D_s} = \frac{1}{G_F m_\ell \left(1 - \frac{m_\ell^2}{M_{D_s}^2}\right) |V_{cs}|} \sqrt{\frac{8\pi \mathcal{B}(D_s \rightarrow \ell \nu)}{M_{D_s} \tau_{D_s}}}$$

Charm: last but not least...


D mixing was discovered at Belle and BaBar...

... and there remains a lot for us to do in the era of super B factories.




Rare τ decays

Example: lepton flavour violating decay $\tau \rightarrow \mu \gamma$




LFV in tau decays: present status

Lepton flavour violation (LFV) in tau decays: would be a clear sign of new physics




LFV and New Physics



- SUSY + Seesaw ($m_{\tilde{l}}^2$)₂₃₍₁₃₎
- Large LFV $Br(\tau \rightarrow \mu \gamma) = O(10^{-7 \sim 9})$


$$Br(\tau \rightarrow \mu \gamma) \approx 10^{-6} \times \left(\frac{(m_{\tilde{L}}^2)_{32}}{\bar{m}_{\tilde{L}}^2} \right) \left(\frac{1 \text{ TeV}}{m_{\text{SUSY}}} \right)^4 \tan^2 \beta$$

$\tau \rightarrow 3l, l \eta$



- Neutral Higgs mediated decay.
- Important when $M_{\text{SUSY}} \gg \text{EW scale}$.

$$Br(\tau \rightarrow 3\mu) = 4 \times 10^{-7} \times \left(\frac{(m_{\tilde{L}}^2)_{32}}{\bar{m}_{\tilde{L}}^2} \right) \left(\frac{\tan \beta}{60} \right)^6 \left(\frac{100 \text{ GeV}}{m_A} \right)^4$$




model	$Br(\tau \rightarrow \mu \gamma)$	$Br(\tau \rightarrow 3l)$
mSUGRA+seesaw	10^{-7}	10^{-9}
SUSY+SO(10)	10^{-8}	10^{-10}
SM+seesaw	10^{-9}	10^{-10}
Non-Universal Z'	10^{-9}	10^{-8}
SUSY+Higgs	10^{-10}	10^{-7}

Integ. Lum. (ab^{-1})


Peter Križan, Ljubljana

New hadrons at B-factories





Coloured boxes: exotic candidates

Molecular states?



Tetra-quarks?



Hybrids?





Resonant substructure in $\Upsilon(5S) \rightarrow h_b(nP) \pi^+ \pi^-$

Look at $M(h_b \pi^+) = MM(\pi^-)$
measure $\Upsilon(5S) \rightarrow h_b \pi \pi$
yield in bins of $MM(\pi)$

Exclusive searches:

Observed in $\Upsilon(5S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$,
 $\Upsilon(2S) \pi^+ \pi^-$ and $\Upsilon(3S) \pi^+ \pi^-$



$Z_b(10610)$ $M = 10608.1 \pm 1.7 \text{ MeV}$
 $\Gamma = 15.5 \pm 2.4 \text{ MeV}$


$Z_b(10650)$ $M = 10653.3 \pm 1.5 \text{ MeV}$
 $\Gamma = 14.0 \pm 2.8 \text{ MeV}$

Seen in 5 different final states,
parameters are consistent

$J^P=1^+$ in agreement with data;
other J^P are disfavored

→ What is the nature of Z_b^+ ? Molecules, tetraquarks, cusps, ... ?


Charged charmonium in $Y(4260) \rightarrow J/\psi \pi^+ \pi^-$



$Y(4260)$ produced via ISR
(Initial State Radiation)

Observed also by BES III.
They also recently found a peak
in $(DD^*)^+$ at 3885 MeV
PRL110, 252001 (2013)
PRL112, 022001 (2014)

Look for a resonance in $J/\psi \pi^+$



Found! $\rightarrow Z_c^+(3895)$

PRL110, 252002 (2013)

very similar to
 $\Upsilon(5S) \rightarrow Z_b^+ \pi^- \rightarrow \Upsilon(1s) \pi^+ \pi^-$

B factories: a success story

- Measurements of CKM matrix elements and angles of the unitarity triangle
- Observation of direct CP violation in B decays
- Measurements of rare decay modes (e.g., $B \rightarrow \tau \nu$, $D \tau \nu$)
- $b \rightarrow s$ transitions: probe for new sources of CPV and constraints from the $b \rightarrow s \gamma$ branching fraction
- Forward-backward asymmetry (A_{FB}) in $b \rightarrow s l^+ l^-$
- Observation of D mixing
- Searches for rare τ decays
- Discovery of exotic hadrons including charged charmonium- and bottomonium-like states

B factories **remain competitive** in many measurements because of their **unique capabilities**.

What next?

Next generation: Super B factories → Looking for NP

→ Need much more data (almost two orders!)


However: it will be a different world in three years, there is a hard competition from LHCb and BESIII

Still, e^+e^- machines running at (or near) $\Upsilon(4s)$ will have considerable advantages in several classes of measurements, and will be complementary in many more

→ Physics at Super B Factory, arXiv:1002.5012 (Belle II)
→ SuperB Progress Reports: Physics, arXiv:1008.1541 (SuperB)




Need $O(100x)$ more data \rightarrow Next generation B-factories

Peak Luminosity Trends (e^+e^- collider)



How to do it?

→ upgrade the existing KEKB and Belle facility



How to increase the luminosity?

$$L = \frac{\gamma_{e\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \left(\frac{I_{e\pm} \xi_{\zeta y}^{e\pm}}{\beta_y^*} \right) \left(\frac{R_L}{R_{\xi_y}} \right)$$

Lorentz factor
 Beam current
 Beam-beam parameter
 Classical electron radius
 Beam size ratio@IP
 1 - 2 % (flat beam)
 Vertical beta function@IP
 Lumi. reduction factor (crossing angle) & Tune shift reduction factor (hour glass effect)
 0.8 - 1 (short bunch)

- (1) Smaller β_y^*
- (2) Increase beam currents
- (3) Increase $\xi_{\zeta y}$

“Nano-Beam” scheme


Collision with very small spot-size beams

Invented by Pantaleo Raimondi for SuperB

How big is a nano-beam ?




How to go from an excellent accelerator with world record performance – KEKB – to a 40x times better, more intense facility?

In KEKB, colliding electron and positron beams were already **much thinner than a human hair...**





... For a 40x increase in intensity you have to make the beam as thin as a **few x100 atomic layers!**

KEKB → SuperKEKB




Replace short dipoles with longer ones (LER)



Add / modify RF systems for higher beam current

Low emittance positrons to inject

Damping ring



Positron source

New positron target / capture section





Low emittance gun

Low emittance electrons to inject

Redesign the lattices of HER & LER to squeeze the emittance

TiN-coated beam pipe with antechambers




To get x40 higher luminosity

Installation of 100 new long LER bending magnets done




Installation of HER wiggler chambers in Oho straight section is done.



Add / modify RF systems for higher beam current

Low emittance positrons to inject



Low emittance gun
Low emittance electrons to inject

Damping ring tunnel: built!



Entirely new LER beam pipe with ante-chamber and Ti-N coating





Beam pipe is made of aluminum.



Fabrication of the LER arc beam pipe section is completed

Al ante-chamber before coating



After TiN coating
before baking

After baking





All 100 4 m long dipole magnets have been successfully installed in the low energy ring (LER)!


Three magnets per day !

Installing the 4 m long LER dipole **over** the 6 m long HER dipole (remains in place).

Magnet installation



field measurement



move into tunnel



carry on an air-pallet

Installation of 100 new LER bending magnets done



carry over existing HER dipole

installation done





Critical issues at $L = 8 \times 10^{35} / \text{cm}^2 / \text{sec}$


- ▶ **Higher background ($\times 10\text{-}20$)**
 - radiation damage and occupancy
 - fake hits and pile-up noise in the EM
- ▶ **Higher event rate ($\times 10$)**
 - higher rate trigger, DAQ and computing
- ▶ **Require special features**
 - low $p \mu$ identification $\leftarrow s \mu \mu$ recon. eff.
 - hermeticity $\leftarrow \nu$ "reconstruction"

Solutions:


- ▶ Replace inner layers of the vertex detector with a pixel detector.
- ▶ Replace inner part of the central tracker with a silicon strip detector.
- ▶ Better particle identification device
- ▶ Replace endcap calorimeter crystals
- ▶ Faster readout electronics and computing system.



Belle II Detector




Belle II Detector (in comparison with Belle)



- SVD: 4 DSSD lyrs → 2 DEPFET lyrs + 4 DSSD lyrs
- CDC: small cell, long lever arm
- ACC+TOF → TOP+A-RICH
- ECL: waveform sampling (+pure CsI for endcaps)
- KLM: RPC → Scintillator +MPPC (endcaps, barrel inner 2 lyrs)



In colours: new components

Belle II Detector – vertex region




Beryllium beam pipe
2cm diameter

Vertex Detector
2 layers DEPFET + 4 layers DSSD




Belle II CDC

Wire Configuration



Much bigger than in Belle!





Wire stringing in a clean room

- thousands of wires,
- 1 year of work...




Aerogel RICH (endcap PID)



Clear Cherenkov image observed

Cherenkov angle distribution




6.6 σ π/K at 4GeV/c !

Peter Križan, Ljubljana

RICH with a novel "focusing" radiator – a two layer radiator

Employ multiple layers with different refractive indices → Cherenkov images from individual layers overlap on the photon detector.




Radiator with multiple refractive indices

How to increase the number of photons without degrading the resolution?

normal

→ stack two tiles with different refractive indices: “focusing” configuration






→ focusing radiator

Such a configuration is only possible with aerogel (a form of Si_xO_y) – material with a tunable refractive index between 1.01 and 1.13.




Focusing configuration – data

Increases the number of photons without degrading the resolution

4cm aerogel single index







2+2cm aerogel




→ NIM A548 (2005) 383

DIRC (@BaBar) - detector of internally reflected Cherenkov light




4 x 1.225 m Bars glued end-to-end


Belle II Barrel PID: Time of propagation (TOP) counter




- Cherenkov ring imaging with **precise time measurement**.
- Uses internal reflection of Cherenkov ring images from quartz like the BaBar DIRC.
- Reconstruct Cherenkov angle from two hit coordinates and the time of propagation of the photon
 - Quartz radiator (2cm thick)
 - **Photon detector (MCP-PMT)**
 - Excellent time resolution ~ 40 ps
 - Single photon sensitivity in 1.5




Barrel PID: Time of propagation (TOP) counter




Example of Cherenkov-photon paths for 2 GeV/c π^\pm and K^\pm .



TOP image



Pattern in the coordinate-time space ('ring') of a pion hitting a quartz bar with ~ 80 MAPMT channels



Time distribution of signals recorded by one of the PMT channels: different for π and K (\sim shifted in time)

The Belle II Collaboration



A very strong group of ~ 600 highly motivated scientists!

SuperKEKB/Belle II Status


Funding

- ~100 MUS for machine approved in 2009 -- Very Advanced Research Support Program (FY2010-2012)
- Full approval by the Japanese government in December 2010; the project was finally in the JFY2011 budget as approved by the Japanese Diet in 2011
- Non-Japanese funding agencies have also allocated sizable funds for the upgrade of the detector.


SuperKEKB and Belle II construction proceeding, nearly on schedule.

Commissioning start delayed 9 months from original plan, now scheduled for October 2015.

SuperKEKB luminosity projection



Summary



- B factories have proven to be an excellent tool for flavour physics, with **reliable long term** operation, constant **improvement** of the performance, **achieving and surpassing** design performance
- Super B factory at KEK under construction 2010-15 → SuperKEKB+Belle II, **L x40, construction at full speed** – the biggest particle physics project under preparation
- Expect a new, exciting era of discoveries, complementary to the LHC



Additional slides


Complementary to LHCb

Observable	Expected th. accuracy	Expected exp. uncertainty	Facility
CKM matrix			
$ V_{us} [K \rightarrow \pi \ell \nu]$	**	0.1%	<i>K</i> -factory
$ V_{cb} [B \rightarrow X_c \ell \nu]$	**	1%	Belle II
$ V_{ub} [B_d \rightarrow \pi \ell \nu]$	*	4%	Belle II
$\sin(2\phi_1) [c\bar{c}K_S^0]$	***	$8 \cdot 10^{-3}$	Belle II/LHCb
ϕ_2		1.5°	Belle II
ϕ_3	***	3°	LHCb
CPV			
$S(B_s \rightarrow \psi\phi)$	**	0.01	LHCb
$S(B_s \rightarrow \phi\phi)$	**	0.05	LHCb
$S(B_d \rightarrow \phi K)$	***	0.05	Belle II/LHCb
$S(B_d \rightarrow \eta' K)$	***	0.02	Belle II
$S(B_d \rightarrow K^*(\rightarrow K_S^0 \pi^0) \gamma)$	***	0.03	Belle II
$S(B_s \rightarrow \phi \gamma)$	***	0.05	LHCb
$S(B_d \rightarrow \rho \gamma)$		0.15	Belle II
A_{SL}^d	***	0.001	LHCb
A_{SL}^s	***	0.001	LHCb
$A_{CP}(B_d \rightarrow s \gamma)$	*	0.005	Belle II
rare decays			
$\mathcal{B}(B \rightarrow \tau \nu)$	**	3%	Belle II
$\mathcal{B}(B \rightarrow D \tau \nu)$		3%	Belle II
$\mathcal{B}(B_d \rightarrow \mu \nu)$	**	6%	Belle II
$\mathcal{B}(B_s \rightarrow \mu \mu)$	***	10%	LHCb
zero of $A_{FB}(B \rightarrow K^* \mu \mu)$	**	0.05	LHCb
$\mathcal{B}(B \rightarrow K^{(*)} \nu \nu)$	***	30%	Belle II
$\mathcal{B}(B \rightarrow s \gamma)$		4%	Belle II
$\mathcal{B}(B_s \rightarrow \gamma \gamma)$		$0.25 \cdot 10^{-6}$	Belle II (with 5 ab^{-1})
$\mathcal{B}(K \rightarrow \pi \nu \nu)$	**	10%	<i>K</i> -factory
$\mathcal{B}(K \rightarrow e \pi \nu) / \mathcal{B}(K \rightarrow \mu \pi \nu)$	***	0.1%	<i>K</i> -factory
charm and τ			
$\mathcal{B}(\tau \rightarrow \mu \gamma)$	***	$3 \cdot 10^{-9}$	Belle II
$ q/p _D$	***	0.03	Belle II
$\arg(q/p)_D$	***	1.5°	Belle II

→ Need both **LHCb** and **super B factories** to cover all aspects of precision flavour physics

B. Golob, KEK FF Workshop, Feb. 2012

$|V_{ub}|$ from $B^0 \rightarrow \pi^- \ell^+ \nu$ exclusive decays



Yield: 2d fit in $M_{bc}=M_{ES}$
and ΔE , bins of q^2

$$m_{bc} = \sqrt{E_{\text{beam}}^2 - |\vec{p}_{\pi} + \vec{p}_{\ell} + \vec{p}_{\nu}|^2}$$

$$\Delta E = E_{\text{beam}} - (E_{\pi} + E_{\ell} + E_{\nu})$$

$$\mathcal{B} = (1.41 \pm 0.05 \pm 0.07) \cdot 10^{-4}$$

untagged

BaBar, PRD83, 032007 (2011)

$$\mathcal{B} = (1.42 \pm 0.05 \pm 0.07) \cdot 10^{-4}$$

untagged loose neutrino

BaBar, PRD83, 052011 (2011)

$$\mathcal{B} = (1.49 \pm 0.04 \pm 0.07) \cdot 10^{-4}$$



untagged

Belle, PRD83, 071101 (2011)

$$\mathcal{B} = (1.49 \pm 0.09 \pm 0.07) \cdot 10^{-4}$$

hadron tag

Belle, PRD88, 032005 (2013)



$|V_{ub}|$ extraction: fit data +
LQCD points in $q^2 = (p_{\ell} + p_{\nu})^2 = (p_B - p_{\pi})^2$

Belle + BaBar +
FNAL/MILC

$$|V_{ub}| = (3.23 \pm 0.30) \cdot 10^{-3}$$



$|V_{ub}|$ from inclusive decays $B \rightarrow X_u \ell^+ \nu$

The other possibility: **inclusive $b \rightarrow u$** measurement by measuring

- lepton spectrum in semileptonic $b \rightarrow u \ell^+ \nu$ decays, or by using
- tagged events (e.g. fully reconstruct one of the B's, and then measure the rate vs mass of the hadronic system X_u)

Inclusive decays

$$|V_{ub}| = (4.42 \pm 0.20 \text{ (exp)} \pm 0.15 \text{ (th)}) \cdot 10^{-3}$$

vs exclusive decays

$$|V_{ub}| = (3.23 \pm 0.30) \cdot 10^{-3}$$

→ Tension between inclusive and exclusive decays is still there - and not understood


Extrapolation to Belle II (1)

Alexander Ermakov (FPCP14):

$|V_{ub}|_{\text{exc}}$ vs $|V_{ub}|_{\text{inc}}$ “tension” is still here after years of experimental and theoretical efforts
Just statistics?

A systematic effect in experiment. or theory or both?


Belle II expected to settle this.



Belle II will reduce the uncertainties on $|V_{ub}|$

But also provide much more consistency checks for theory and experimental effects

$|V_{ub}|$ extrapolation for Belle II (2)



	Statistical	Systematic	Total Exp	Theory	Total
	(reducible, irreducible)				
$ V_{ub} $ exclusive (had. tagged)					
711 fb ⁻¹	3.0	(2.3, 1.0)	3.8	8.7 (2.0)	9.5 (4.3)
5 ab ⁻¹	1.1	(0.9, 1.0)	1.7	4.0 (2.0)	4.4 (2.6)
50 ab ⁻¹	0.4	(0.3, 1.0)	1.1	2.0	2.3
$ V_{ub} $ exclusive (untagged)					
605 fb ⁻¹	1.4	(2.1, 0.8)	2.9	8.7 (2.0)	9.1 (4.0)
5 ab ⁻¹	0.5	(0.8, 0.8)	1.2	4.0 (2.0)	4.2 (2.4)
50 ab ⁻¹	0.2	(0.3, 0.8)	0.9	2.0	2.2
$ V_{ub} $ inclusive					
605 fb ⁻¹ (old <i>B</i> tag)	4.5	(3.7, 1.6)	6.0	2.5–4.5	6.5–7.5
5 ab ⁻¹	1.1	(1.3, 1.6)	2.3	2.5–4.5	3.4–5.1
50 ab ⁻¹	0.4	(0.4, 1.6)	1.7	2.5–4.5	3.0–4.8

Assumption is theory error down to 2% for exclusive and 2-4 % for inclusive modes

Most promising are exclusive analysis with hadronic tags: to perform clean and detailed exploration of exclusive $b \rightarrow u$ modes spectra. Improvements on theory predictions need as well ($B \rightarrow \rho l \nu$ lattice)

Untagged analyses still competitive for $|V_{ub}|$ measurement

$B \rightarrow D^* l \nu$ and $B \rightarrow D l \nu$

	Statistical	Systematic	Total Exp	Theory	Total
	(reducible, irreducible)				
$ V_{cb} $ exclusive : F(1)					
711 fb ⁻¹	0.6	(2.8, 1.1)	3.1	1.8	3.6
5 ab ⁻¹	0.2	(1.1, 1.1)	1.5	1.0	1.8
50 ab ⁻¹	0.1	(0.3, 1.1)	1.2	0.8*	1.4
$ V_{cb} $ exclusive : G(1)					
423 fb ⁻¹	4.5	(3.1, 1.2)	5.6	2.2	3.6
5 ab ⁻¹	1.3	(0.9, 1.2)	2.0	1.5*	2.7
50 ab ⁻¹	0.6	(0.4, 1.2)	1.4	1.0*	1.7

Similar level of accuracy from $B \rightarrow D^* l \nu$ and $B \rightarrow D l \nu$

$B \rightarrow X_c l \nu$ inclusive at Belle II

(Modest) improvement of experimental uncertainties expected.


- Better determination of $B \rightarrow D^{**} l \nu$ component
- Improved control on the tag B normalization
- Largest experimental sys effect from PID and tracking

We expect a 0.5% ultimate systematic uncertainty

We assume theory uncertainty at 1% that will saturate the error budget

Detailed exploration of $B \rightarrow D \pi l \nu$

Hopefully will solve “puzzles” like the gap between inclusive and exclusive rates (and V_{cb} tensions)




Fitted $D^{(*)} \pi$ mass spectrum of
Phys.Rev.Lett. 101 (2008) 261802

ϕ_3 : Binned Dalitz plot analysis

Solution: use binned Dalitz plot and deal with numbers of events in bins.

[A. Giri, Yu. Grossman, A. Soffer, J. Zupan, PRD 68, 054018 (2003)]

[A. Bondar, A. P. EPJ C 47, 347 (2006); EPJ C 55, 51 (2008)]



$$M_i^\pm = h\{K_i + r_B^2 K_{-i} + 2\sqrt{K_i K_{-i}}(x_\pm c_i + y_\pm s_i)\}$$

$$x_\pm = r_B \cos(\delta_B \pm \phi_3) \quad y_\pm = r_B \sin(\delta_B \pm \phi_3)$$

M_i^\pm : numbers of events in $D \rightarrow K_S^0 \pi^+ \pi^-$ bins from $B^\pm \rightarrow DK^\pm$

K_i : numbers of events in bins of flavor $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ from $D^* \rightarrow D\pi$.

c_i, s_i contain information about strong phase difference between symmetric

Dalitz plot points $(m_{K_S^0 \pi^+}^2, m_{K_S^0 \pi^-}^2)$ and $(m_{K_S^0 \pi^-}^2, m_{K_S^0 \pi^+}^2)$:

$$c_i = \langle \cos \Delta\delta_D \rangle, \quad s_i = \langle \sin \Delta\delta_D \rangle$$

ϕ_3 : Obtaining c_i, s_i

Coefficients c_i, s_i can be obtained in $\psi(3770) \rightarrow D^0 \bar{D}^0$ decays.
Use quantum correlations between D^0 and \bar{D}^0 .

- If both D decay to $K_S^0 \pi^+ \pi^-$, the number of events in i -th bin of $D_1 \rightarrow K_S^0 \pi^+ \pi^-$ and j -th bin of $D_2 \rightarrow K_S^0 \pi^+ \pi^-$ is

$$M_{ij} = K_i K_{-j} + K_{-i} K_j - 2\sqrt{K_i K_{-i} K_j K_{-j}}(c_i c_j + s_i s_j).$$

\Rightarrow constrain c_i and s_i .

- If one D decays to a CP eigenstate, the number of events in i -th bin of another $D \rightarrow K_S^0 \pi^+ \pi^-$ is

$$M_i = K_i + K_{-i} \pm 2\sqrt{K_i K_{-i}} c_i.$$


\Rightarrow constrain c_i .

c_i, s_i measurement has been done by CLEO and can be done in future at BES-III.




$\sin 2\phi_1 (= \sin 2\beta)$ vs. $\mathcal{B}(B \rightarrow \tau\nu)$

Tension between $\mathcal{B}(B \rightarrow \tau\nu)$ and $\sin 2\phi_1$
very much reduced (from $\sim 2.5 \sigma$)




Extracting ϕ_2 : isospin relations


$$B^0 \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$$



$$T \sim V_{ub}^* V_{ud} \sim \lambda^3$$



$$P \sim V_{tb}^* V_{td} \sim \lambda^3$$




$$T_c \sim V_{ub}^* V_{ud}$$

No penguin!

Constraint: relation of decay amplitudes in the SU(2) symmetry

$$A^{+0} = \frac{1}{\sqrt{2}} \bar{A}^{+-} + \bar{A}^{00}$$

$$A^{-0} = \frac{1}{\sqrt{2}} A^{+-} + A^{00}$$




ϕ_2 $B^0 \rightarrow \rho^0 \rho^0$ 6D fit to $\Delta E, \mathcal{F}_{S/B}, m_1(\pi^+\pi^-), m_2(\pi^+\pi^-), \cos\theta_{\text{Hel}}^1, \cos\theta_{\text{Hel}}^2$

PRD 89 072008, (2014)



$$\mathcal{B}(B^0 \rightarrow \rho^0 \rho^0) = (1.02 \pm 0.30 \pm 0.15) \times 10^{-6}$$

3.4 σ significance

$$f_L = 0.21_{-0.22}^{+0.18} \pm 0.15$$

 $(f_L: 2.1\sigma \text{ tension with Babar's result})$ 

Signal enhanced projections



ϕ_2 $B \rightarrow \rho\rho$ ϕ_2 from isospin analysis (longitudinal polarization)

- constraints using Belle results only

$$\phi_2 = (84.9 \pm 13.5)^\circ$$

$$\Delta\phi_2 = (0.0 \pm 10.4)^\circ$$

⇒ Belle needs updates on


- $B^0 \rightarrow \rho^+ \rho^-$

B, f_L from 275M $B\bar{B}$ (PRL96, 171801 (2006))

\mathcal{CP} from 535M $B\bar{B}$ (PRD76, 011104 (2007))

- $B^\pm \rightarrow \rho^\pm \rho^0$

85M $B\bar{B}$ (PRL91, 221801 (2006))




$$\text{W.A.: } \phi_2 = (89.9^{+5.4}_{-5.3})^\circ$$

[prospect for Belle2 $\phi_2^{\rho\rho} = (X \pm 3)^\circ$]

$\pi^+ \pi^- S_{CP}$ vs C_{CP}

HFAAG
Moriond 2014
PRELIMINARY



$$(\mathcal{A}_{CP} = -\mathcal{C}_{CP})$$

measurements in good agreement,

current average:

- $\mathcal{S}_{CP}^{\pi^+\pi^-} = -0.66 \pm 0.06$
- $\mathcal{A}_{CP}^{\pi^+\pi^-} = +0.31 \pm 0.05$

significant amount of penguin
contribution

Conclusions

$B^0 \rightarrow \omega K_S^0$: BR and time dependent CPV

- ▶ $\mathcal{S}_{\omega K_S^0} = +0.91 \pm 0.32 \pm 0.05$
- ▶ first evidence (3.1σ) for CPV
- ▶ four out of five parameters world's most precise results

$B^0 \rightarrow \eta' K^0$: time dependent CPV

- ▶ $\mathcal{S}_{\eta' K^0} = +0.68 \pm 0.07 \pm 0.03$
- ▶ Most precise determination of CPV parameters

$B^0 \rightarrow K_S^0 \eta \gamma$: time dependent CPV


- ▶ no significant CPV observed

So far, everything is consistent with SM

- ▶ most analyses are statistically limited

$$\sin(2\beta^{\text{eff}}) \equiv \sin(2\phi_1^{\text{eff}}) \quad \text{HFAG}$$

Moriond 2014
PRELIMINARY



Naïve $b \rightarrow s q \bar{q}$ average 0.655 ± 0.032